AGRI-VETA LTD

AGRI VETA

VETERENARY MEDICINE AND VITAMINS

Agri Veta upholds the highest standards of quality and integrity, ensuring that all our veterinary medicines and vitamin products meet the needs of veterinarians, livestock growers, and pet owners alike.



Agriveta VitaBoost ADEB

AD3E + B Complex Injection

COMPOSITION:

Each ml contains:

Vitamine A.....2000 IU
Vitamine D3.....2000 IU
Vitamine E.....0.5 mg
Vitamine B1.....0.3 mg
Vitamine B.....0.06 mg
Vitamine B.....0.9 mg
Niacinamide.....3 mg
Folic acid.....0.3 mg
Excipients (ad.).....1 ml

INDICATIONS: This product is indicated in the prevention and treatment of vitamin deficiencies in cattle, sheep and goats. Particularly during periods of illness, convalescence and general unthriftiness. Improvement of feed convertion.

ADMINISTRATION AND DOSAGE: 1 ml/10 kg body weight by SC, IM or slow IV.

CONTRAINDICATIONS:

This product is contraindicated in animals with known hypersensitivity to any of the ingredients.

SIDE EFFECTS: No undesirable effects are to be expected when the prescribed dosage regimen is followed.

WITHDRAWAL PERIOD:

Not required.

AGRI-VETA LTD

Agriveta PainRelief-C

Analgin + Vitamin C Injection 30% + 10%

COMPOSITION:

Each ml contains:

Vanalgin.....300 mg

Vitamin C.....100 mg

Excipients .....1 ml

INDICATIONS: Antipyretic, anti-inflammatory, antistress, relieves pain after injuries, surgery or other acute pain, Analgin - C should be used in combination with antibiotic for better treatment of infectious diseases: infections, arthritis dislocated and sprained in cattle and poultry.

CONTRAINDICATIONS: If you are allergic to Analgin and other pyrazolone derivatives. Animals sensitives to any ingredients in drug.

ADMINISTRATION AND DOSAGE: Cattle, camels: 1 ml/15 kg body weight/day. Goats, sheep, dogs, cats: 1 ml/ 8-10 kg body weight/day. Poultry: 1 ml/5-6 kg body weight/day or 5 ml/2-3 liters of water/day. By Intramuscular (I.M.) or Intravenous (slowly) injection. Twice a day until signs of symptom disappear.

AGRI-VETA LTD

Agriveta Cardioboost

AMP5 + VB12+ Salbutamol Injection

COMPOSITION:

Each ml contains:

Adenosine-5-monophosphate..... 200 mg

Salbutamol.....2.5 mg

Vitamin B1.....12 mcg

Excipients.....1ml

INDICATIONS: Production of coronary vasodilation in anticipation of severe cardiac load in athletic animals or in treatment of myocardial oxygen deficit (as evidenced by performance linked ECG changes). Muscle cramp, 'tying up' prevention. salbutamol sulphate is indicated for the relief of severe bronchospasm associated with asthma or bronchitis.

PRESENTATION: 20 ml sterile glass multi-dose vial.

ADMINISTRATION AND DOSAGE: Administer by intramuscular injection only. Give two injections: one 24 hours before and again four to six hours before strenuous exercise.

Horses: 2.5ml/100 kg.

Dogs: 0.5 ml/10 kg.

AGRI-VETA LTD

Agriveta TrypCure Plus

OXYTETRACYCLINE HCL, DIACETURATE DIMINAZENE, ANTIPYRINE AND CYANOCOBALAMIN.

COMPOSITION:

Each ml contains:

Oxytetracycline Hydrochloride.....100 mg

Diminazene Diaceturate.....3.5 mg

Antipyr.....150 mg

Cyanocobalamin.....50. µg

Excipients: (Magnesium chloride hexahydrated, propylene glycol USP, monoethanolamine , EDTA disodium, sodium formaldehyde sulfoxylate, 2-pyrrolidone, P.V.K-17 (Kollidon K-17), water for injectable)..... ml.

INDICATIONS: In cattle, equines, sheep, goats and canines for the treatment of diseases caused by:) Anaplasma sp., Babesia bigemina, B. bovis, B. ovis, B. canis, Theileria equi, Trypanosoma vivaxy T. evansi.

DESCRIPTION: Translucid homogeneous solution, free of strange visible material.

WITHHOLDING PERIOD: Treated animals can't be slaughtered for human consumption before twen-ty-one (21) days after treatment. Milk produced during the treatment or before six (6) days after treatment can't be use for human consumption.

PRODUCT FLAMMABILITY: It can be flammable to high temperatures. It is not explosive because of heat, mechanical impact or static discharge.

AGRI-VETA LTD

Agriveta Histaclear

Chlorphenamine maleate Injection 1%

COMPOSITION:
Each ml contains:
Chlorphenamine maleate.....10 mg
Excipients .....1 ml

INDICATIONS: For the symptomatic treatment of conditions associated with histamine release. Such as urticaria, drug allergy, allergic dermatitis, serum sickness, angioedema and other allergic reactions of the skin and mucous membranes with congestion, edema and itching.It also has a certain curative effect on gastrointestinal recovery and diarrhea caused by allergies. It can also be used for diseases associated with histamine release due to tissue damage, such as burns, frostbite, eczema, septic metritis, etc.

ADMINISTRATION AND DOSAGE

Intramuscular or intravenous use. Intravenous injection should be slow and, if necessary, discontinued for a few minutes. Adult animals: 0.5 mg Chlorphenamine maleate/kg body weight (equivalent to 5 ml/100 kg body weight), once a day for three consecutive days. Calves: 1 mg Chlorphenamine maleate / kg body weight (equivalent to 10 ml / 100 kg body weight), once a day for three consecutive days.

CONTRAINDICATIONS: Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

SIDE EFFECTS: Chlorphenamine has a weak sedative effect.

WITHDRAWAL PERIOD
Meat and offal: 1 day.
Milk: 12 hours.

AGRI-VETA LTD

Agriveta MagCal Forte

Minerals

COMPOSITION: Each ml contains: Magnesium gluconate.....15 g
Magnesium chloride, hexahydrate.....5g
Zinc sulfate.....0.5 g
Calcium hypophosphite.....2.13 g
Magnesium hypophosphite.....3 g
Boric acid.....3,5 g
Formulation agents.....q.s.

INDICATIONS: Treatment and prevention of magnesium deficiencies. It is administered preventively in herds of flocks with a previous history of hypoenergetic syndromes and/or hypomagnesemia. Its application is recommended in situations of severe stress (transport, weaning, storms, consumption of winter pastures, shearing, start of the last third period of gestation in sheep). Treatment of metabolic dysfunctions caused by a sudden alteration of the energy and/or magnesium requirements in ruminants (grass tetany, ketosis in dairy cows, fat cow syndrome, pregnancy toxemia in sheep, hypoenergetic, syndrome in sheep).

ANIMAL SPECIES TO WHICH IT IS INTENDED; Cattle and sheep.

DOSAGE: Cattle weighing up to 500 k.l.w.: 20 ml subcutaneously; 300 to 500 ml intravenously in a slow way. Steers: 15 ml subcutaneously: 100 ml intravenously. Calves: 5 to 10 ml subcutaneoulsy: 50 ml intravenously. Sheep: 5 ml subcutaneously; 20 to 50 ml intravenously.

AGRI-VETA LTD

Agriveta TheilGuard

Buparvaquone Injection 5%

COMPOSITION: Each ml contains:

Buparvaquone:.....50 mg

Excipients (ad.).....1 ml

INDICATIONS: For treatment of tick-transmitted theileriosis caused by the intracellular protozoan parasites Theileria parva (East Coast fever, Corridor disease, Zimbabwean theileriosis) and T. annulata (tropical theileriosis) in cattle. It is active against both the schizont and piroplasm stages of Theileria spp. and may be used during the incubation period of the disease, or when clinical signs are apparent

ADMINISTRATION AND DOSAGE: IFor intramuscular injection. The general dosage is 1 ml per 20 kg body weight. It is recommended to be given into the muscles of the neck. In severe cases the treatment may be repeated within 48-72 hours. Do not administer more than 10 ml per injection site. Successive injections should be administered at different sites.

CONTRAINDICATIONS: Due to the inhibiting effects of theileriosis on the immune system, vaccination should be delayed until the animal has recovered from theileriosis.

WITHDRAWAL PERIOD:

Meat: 42 days. Milk: 48 hours.

SHELF LIFE: Shelf life of the veterinary medicinal product as packaged for sale: 36 months. Shelf life after first opening the immediate packaging: 28 days.

AGRI-VETA LTD

Agriveta Cortiplex

Injectable solution

COMPOSITION: Each ml contains:

Prednisolone.....1 mg
Hydrocortisone.....50 mg
Phenyle Alanine.....100 mg
Isoleucin.....50 mg
Excipients (ad.).....1 ml

INDICATIONS: Musculoskeletal Conditions. As with other adrenal steroids, Corta Predin super Sterile Aqueous Suspension has been found useful in alleviating the pain and lameness associated with acute localized arthritic conditions and generalized arthritic conditions. Allergic Conditions. This preparation is especially beneficial in relieving pruritus and inflammation of allergic dermatitis, acute moist dermatitis, dry eczema, urticaria, bronchial asthma, pollen sensitivities and otitis externa in dogs; allergic dermatitis and moist and dry eczema in cats. Corta Predin super may be lifesaving. acting to inhibit the inflammatory reaction.

ADMINISTRATION AND DOSAGE: Following intramuscular injection of methylprednisolone ace- tate, a prolonged systemic effect results. Dogs and Cats: The average intramuscular dose for dogs is 20 mg. The average intramuscular dose for cats is 10 mg with a range up to 20 mg. Horses:The usual intramuscular dose for horses is 200 mg repeated as necessary.

AGRI-VETA LTD

Agriveta CuB12 Tonic

Copper + VB12 Injection 2% +0.1%

COMPOSITION:

Each ml contains:

Copper (as copper methionat) .....20 mg

Vitamin B12.....1 mg

Excipients .....1 ml

INDICATIONS: Treatment of hypocuprosis in cattle with concurrent anaemia. Prevention and treatment of copper deficiency in sheep.

ADMINISTRATION AND DOSAGE: Administration is by deep intramuscular injection into the neck area. The dosage and frequency of therapy required depends on the clinical condition of the animal and the copper status, as assessed by blood and liver levels both before and after therapy. The recommended dosage rate is 20 mg copper per 50 kg body weight. Cattle: (Adult): 4-6 ml. - Ewes: 2 ml.

CONTRAINDICATIONS: Do not use in animals with known hypersensitivity to the active ingredient. Do not administration by the intravenous route.

WITHDRAWAL PERIOD:

Meat: Animals must not be slaughtered for human consumption within 21 days of administration.

Milk: Nil.

AGRI-VETA LTD

Agriveta FlexiThera

Phenylbutazone + Dexamethasone Injection

COMPOSITION:

Each ml contains:

Phenylbutazone.....180 mg

Dexamethasone.....0.35 mg

Excipients .....1 ml

DESCRIPTION: This product works as potent analgesic anti-inflammatory for training, specially designed for horses and camels of high performance competition. Dexaphenyl principal actives are dexamethasone and phenylbutazone. Phenylbutazone works as a powerful analgesic and antipyretic which adds effects and strengthens each other with the dexamethasone, generating tolerances that allow to improve the high performance training.

INDICATIONS: Anti-inflammation, analgesic, antipyretic. The combination between of phenylbuta- zone and dexamethasone enhance animal's resistance during training and exercise or working with high pressure.

ADMINISTRATION AND DOSAGE: Apply 5 ml for 5 days - IM (suspend 7 days before the race). IV and IM administration. Horses and camels: 4 ml/ 100 kg BW. Dogs: 0.5 ml/10 kg BW.

CONTRAINDICATION: Do not use for hypersensitive animals to any component.

WITHDRAWAL PERIOD:

7days before slaughter.

AGRI-VETA LTD

Agriveta Flunivet

Flunixin Meglumine Injection 50 mg/ml

COMPOSITION: Each ml contains:
Flunixin (as flunixin meglumine).....50 mg
Excipients .....1 ml

INDICATIONS: Cattle: It is indicated as an adjunct to antimicrobial therapy to reduce clinical signs of acute inflammation in cases of infectious respiratory disease. Horses: It is indicated for the alleviation of inflammatory signs associated with musculoskeletal dis-orders and for the alleviation of visceral pain associated with colic.

ADMINISTRATION AND DOSAGE: For intramuscular or intravenous administration: Cattle: 2 ml per 45 kg body weight. Repeat if necessary at 24 h intervals for up to 3 consecutive days. Horses: 1 ml per 45 kg body weight. Treatment may be repeated once or twice if colic recurs.

WITHDRAWAL PERIOD:
For meat: Cattle: 14 days.
Horses: 28 days.
For milk: Cattle: 2 days.

AGRI-VETA LTD

Agriveta HepaCare

PROTECT LIVER

COMPOSITION:

Acetylmethionine.....45 mg
DL-methionine.....15 mg
Sorbitol.....210 mg
Cocarboxylase.....3 mg
Excipients .....1 ml

INDICATIONS: Hepatoprotective and antitoxic. Protects and stimulates liver function. Metabolic disorders, anemias and states of exhaustion.

ADMINISTRATION AND DOSAGE: Equines and cattle: 50 to 100 ml via IV. Sheep: 5 to 10 ml via IV or IM. - Canines and felines: 1 ml every 10 kpv via IV or IM.

PROPERTIES: Hepa Care is a detoxifier and liver protector. Its different components are enhanced when combined and make Hepa Super a first-choice medication when it comes to protecting and promoting liver function. Acetylmethionine and DL-Methionine are essential amino acids that have liporegulatory activity and constitute a source of methyl for the formation of choline. They protect the liver from poisoning by heavy metals and other cellular toxins and stimulate erythropoiesis in the bone marrow, Sorbitol is a sugar alcohol that acts as a restorative of the liver cell and is a natural source of energy.

Cocarboxylase or Thiamine pyrophosphate is the biologically active form of vitamin B1 (Thiamin). It helps maintain the necessary energy through the production of Adenosine triphosphate (ATP) for cellular functions to be carried out (protein biosynthesis, muscle contraction, etc.), and thus maintain the proper functioning of tissues. Cocarboxylase in the Krebs cycle acts constantly stimulating energy production and its administration reactivates it immediately.

AGRI-VETA LTD

Agriveta FerroPlus

Iron Dextran + VB12 Injection 20% +0.02%

COMPOSITION: Each ml contains:

Piron (as iron dextran).....200 mg
Vitamin B12, cyanocobalamin.....200 mcg
Excipients .....1 ml

DESCRIPTION: Iron dextran is used for prophylaxis and treatment of by iron deficiency caused anaemia in animals. Parenteral administration of iron has the advantage that the necessary amount of iron can be admin -istered in one single dosage. Cyanocobalamin is used for prophylaxis and treat cyanocobalamin deficiency caused anaemia.

INDICATIONS: Prophylaxis and treatment of anaemia in camels, calves, sheep and goats.

ADMINISTRATION AND DOSAGE:

Dosage and method of use of Parkers Endofer needles for calves, sheep, goats and camels: Used by intramuscular injection.
Dosage in one-year-old camels: 5 ml per animal.
Dosage for adult camels: 10 ml per animal. Dosage in calves: 5 ml per animal.
Dosage in sheep and goats: 2 ml per animal.

AGRI-VETA LTD

Agriveta Doramax

INJECTABLE SOLUTION FOR CATTLE AND SHEEP

COMPOSITION: Each ml contains:
Doramectin.....10 mg
Excipients .....1 ml

INDICATIONS: Treatment and control of cattle and sheep parasites sensitive to doramectin.

ADMINISTRATION AND DOSAGE: Subcutaneous or intramuscular single injection at a dosage of 200 ug doramectin / kg (1 ml/50 kg) body weight.

WITHDRAWAL PERIOD: Meat and offals: 35 days. Milk: Do not use in lactating animals producing milk for human consumption.

CONTRAINDICATIONS: Do not use in dogs, as severe adverse reactions may occur. In common with other avermectins, certain breeds of dog, such as collies, are especially sensitive to doramectin and particular care should be taken to avoid accidental consumption of the product. Do not use in case of hypersensitivity to the active substance or any of the excipients.

AGRI-VETA LTD

Agriveta OxyFlu LA

Oxytetracycline + Flunixin Injection 20%+2%

COMPOSITION:

Oxytetracycline.....200 mg
Flunixin (as meglumine).....20 mg
Excipients .....1 ml

INDICATIONS: This product can be used in Treatment of respiration infections and inflammation especially caused by pasteurella spp, acute mastitis caused by Staphylococcus spp and Coli-forms, also for used in treatment of metritis and gastrointestinal infections.

ADMINISTRATION AND DOSAGE: For intramuscular administration. Cattle, sheep and goats: 1 ml per 20 kg body weight. Once a day for 3-5 days. Maximum volume administered per injection site 20 ml.

WITHDRAWAL PERIOD: Meat and offal: 28 days. Not for use in cattle producing milk for human consumption.

AGRI-VETA LTD

Agriveta CortiDex Plus

COMPOSITION:

Each ml contains:

Dexamethasone.....0.1 mg
Hydrocortisone.....50 mg
Phenyle Alanine.....100 mg
Sodium Selenate.....100 mg
Vitamin B12.....200 mcg
Excipients .....1 ml

INDICATIONS: Dexamethasone is a potent glucocorticoid with minimal mineralocorticoid activity. Dexamethasone has ten to twenty times the anti-inflammatory activity of prednisolone. Following intramuscular injection this soluble ester of dexamethasone is rapidly absorbed and hydrolysed to the parent alcohol giving a prompt response which is maintained for approximately 48 hours. This product may be used whenever a parental corticosteroid preparation giving a medium duration of activity is indicated. It can be used as an anti-inflammatory and anti-allergic agent in horses, dogs and cats. The product can also be used as supportive therapy in cases of shock in horses.

ADMINISTRATION AND DOSAGE: This product may be administered by intravenous, intramuscular or subcutaneous injection in horses, dogs and cats. The product may also be given by intra-articular injection in horses. Normal aseptic techniques should be observed. To measure small volumes of less than 1 ml a suitably graduated syringe should be used to ensure accurate administration of the correct dose.

AGRI-VETA LTD

Agriveta HydroCort

Hydrocortisone Injection 5%

COMPOSITION:

Each ml contains:

Hydrocortisone.....50 mg

Excipients .....1 ml

INDICATIONS

It is indicated for severe acute infection or inflammation, bovine ketonemia and sheep pregnancy toxemia, etc.

ADMINISTRATION AND DOSAGE:

For intravenous administration for single dose:

Horses and camels: 4 - 10 ml.

Cattle: 4-10 ml.

Sheep: 0.4 1.5 ml.

WITHDRAWAL PERIOD:

None.

AGRI-VETA LTD

Agriveta Babecare

Imidocarb Dipropionate Injection 12%

COMPOSITION:

Each ml contains:

Imidocarb dipropiona.....120 mg

Excipients .....1 ml

INDICATIONS: It is indicated for treatment and prophylaxis of babesiosis in cattle, for treatment of babesiosis and anaplasmosis in sheep, for treatment of babesiosis in horses and dogs and for treatment of anaplas-mosis in cattle.

ADMINISTRATION AND DOSAGE: For parenteral administration. Calves and cattle: Babesiosis: Treatment of babesiosis: 1.0 ml per 100 kg body weight, subcutaneously. Prevention of babesiosis: 2.5 ml per 100 kg body weight, subcutaneously, one month before exposure. Anaplasmosis: Treatment of anaplasmosis: 2.5 ml per 100 kg bodyweight, subcutaneously. Elimination of the carrier state: 4.0 ml per 100 kg bodyweight, subcutaneously, administered twice with a 14-day interval. Hor ses: Treatment of Babesia caballi: 2.0 ml per 100 kg body weight, once daily for 2 consecutive days, intramuscularly. Treatment of Babesia equi: 3.5 ml per 100 kg body weight, administered 4 times with a 72- hour interval. Sheep: 1.0 ml per 100 kg body weight, intramuscularly. Dogs: 0.5 ml per 10 kg body weight, subcutaneously or intramuscularly. Repeat the dose in 2 weeks, for a total of 2 treatments.

AGRI-VETA LTD

Agriveta WormShield Plus

Ivermectin + Clorsulon Injection 1% + 10%

COMPOSITION: Each ml contains:

Ivermectin.....10 mg
Clorsulon.....100 mg
Excipients (ad.).....1 ml

INDICATIONS: Treatment of gastrointestinal roundworms (adults and fourth-stage larvae), lungworms (adults and fourth-stage larvae), liver fluke (Fasciola hepatica and F. gigantica: adult stages), eye worms, warbles (parasitic stages), sucking lice and mange mites (scabies) in cattle and sheep.

ADMINISTRATION AND DOSAGE: This product should be given only by subcutaneous injection at the recommended dosage level of 1 mL per 50 kg body weight. (based on a dosage level of 200 mcg ivermectin plus 2 mg clorsulon per kg bodyweight) under the loose skin in front of, or behind, the shoulder. Divide doses greater than 10 ml between two injection sites. A sterile 17 gauge 1/2 inch (15-20 mm) needle is recommended. Replace with a fresh sterile needle after every 10-12 animals or sooner if the needle becomes soiled.

CONTRAINDICATIONS: Do not use in animals known to be hypersensitive to the active substance. Do not use by the intravenous or intramuscular route. Do not use in animals producing milk for human consumption.

AGRI-VETA LTD

Agriveta WormShield

Ivermectin 1%

COMPOSITION: Each ml contains:

Ivermectin.....10 mg

Excipients .....1 ml

INDICATIONS: For the treatment and control of gastrointestinal roundworms, lungworms, lice and mange mites in cattle, calves and sheep.

ADMINISTRATION AND DOSAGE: This product should be given only by subcutaneous injection at the recommended dosage level of 1 ml per 50 kg body weight under the loose skin in front of, or behind, the shoulder in cattle, calves and in the neck in sheep. The injection may be given with any standard automatic or single-dose or hypodermic syringe. Use of 17 gauge x 1/2 inch needle is suggested. Replace with a fresh sterile needle after every 10 to 12 animals. Injection of wet or dirty animals is not recommended.

WITHDRAWAL PERIOD;

For meat and offals: Cattle: 49 days. Sheep: 42 days. Do not use in calves to be processed for veal.

SHELF LIFE: Shelf life of the veterinary medicinal product as packaged for sale: 36 months. Shelf life after first opening the immediate packaging: 28 days

AGRI-VETA LTD

Agriveta LinSpect

Lincomycin HCL + Spectinomycin HCL Injection

COMPOSITION:

Each ml contains:

Lincomycin (as Lincomycin Hydrochloride).....50 mg

Spectinomycin (as Spectinomycin Hydrochloride).....100 mg

Excipients .....1 ml

INDICATIONS: Gastrointestinal and respiratory infections caused by lincomycin and spectinomycin sensitive micro-organisms, like Campylobacter, E. coli, Mycoplasma, Salmonella, Staphylococcus, Streptococcus, and Treponema spp., in calves, sheep, goats, cats, dogs and poultr y.

ADMINISTRATION AND DOSAGE: By intramuscular or subcutaneous (poultry) injection. Calves: 1 ml per 10 kg body weight twice on the first day, to be followed by once daily for 2 - 4 days.
Sheep and Goats: 1 ml per 10 kg body weight daily for 3 days. Cats and Dogs: 1 ml per 5 kg body weight, the dosage may be repeated every 12 or 24 hours for maximal 21 days. Poultry: 1 ml per 5 kg body weight by subcutaneous injection daily for 3 days.

WITHDRAWAL PERIODS:

For meat: Calves, goats and sheep: 14 days.
Poultry: 7 days. For milk: 3 days.

AGRI-VETA LTD

Agriveta Methaboost

Methionine Injection 20%

COMPOSITION: Each ml contains:

Methionine.....200 mg

Excipients .....1 ml

DESCRIPTION: Methionine Injection 20% is approved for use in all animal species. Particularly for use in monogastrics, it ensures the adequate supply of sulfur amino acids, especially of the essential methionine. Because of methionine being first-limiting in typical poultry diets, its use has been well established for many years.

INDICATIONS: Methionine Injection 20% is indicated for the prevention or treatment of methionine deficiencies in horses, cattle, calves, sheep, goats, dogs and poultry.

ADMINISTRATION AND DOSAGE: By slow IM or IV injection in the following dose: Horses: 50 ml. Cattle: 50-100 ml. Sheep and Goats: 20 ml. Dogs: up to 10 ml. Repeat after one weak or as directed by the veterinarian. Intramuscular injection is sometimes slightly painful. Required dose can be diluted with water for injection in the amount twice the volume of injection dose.

WITHDRAWAL PERIOD:

None

AGRI-VETA LTD

Agriveta Phenbut

Phenylbutazone Injection 20%

COMPOSITION: Each ml contains:

Phenylbutazone.....200 mg

Excipients .....1 ml

INDICATIONS: arthritis, bursitis, myositis, neuritis,)-Peri) tendinitis and tendovaginitis. Birth trauma, impo-tentia coeundi of bull, muscle injuries and painful injuries like contusions, distortions, haemorrhages and luxa-tions in calves, cattle, goats and sheep.

CONTRA -INDICATIONS: Administration to animals with an impaired liver, renal and/or heart function. Administration to animals with gastric or intestinal ulcers. Administration to very young animals. Caution is necessary when phenylbut 20 is administered together with other strong plasma protein binding drug, which can drive phenylbutazone out of the protein binding (like salicylates, sulphonamides)

ADMINISTRATION AND DOSAGE: For intramuscular or slow intravenous administration. Cattle: 1 ml per 20 kg body weight. Calves, goats, sheep: 1 ml per 10 kg body weight.injection dose.

WITHDRAWAL PERIOD:

For meat: 12 days.

For milk: 4 days.

AGRI-VETA LTD

Agriveta Oxyfen LA

Oxytetracycline + Ketoprofen Injection 20%+3%

COMPOSITION: Each ml contains:

Oxytetracycline.....200 mg

Ketoprofen.....30 mg

Excipients .....1 ml

INDICATIONS:

For the treatment and prevention of all infectious processes that occur with inflammatory reaction, pain and fever. Systemic and local infections, bronchopneumo- nia, mastitis, bacterial enteritis, arthritis, osteoarthritis, urinary tract infection, prostatitis, pododermatitis, keratoconjunctivitis, and others. Nonspecific bacterial and secondary to viral diseases infections

ADMINISTRATION AND DOSAGE:

It may be administered through deep or subcutaneous intramuscular route. 1 ml/kg body weight (20 mg/kg oxytetracycline and 3 mg/kg ketoprophene) as single dose. Although due to product characteristics only one application is required in most part of the cases, some cases may require a second application after 48 hours. it persists, it is convenient to reevaluate the diagnosis.

In intramuscular and subcutaneous applications, the volume must be adjusted to the age and size of the animal so that younger animals will be injected from 1 to 4 ml only, depending on the place of injection. No more than 10 ml per place of injection must be applied in adult cattle, sheep and adult camelids. Each injection requires a sterile needle and syringe which shall be completely clean (they must be sterilized by boiling them in water for 15 minutes). Before taking the solution out from the vial, disinfect the rim of the bottle with an adequate disinfectant such as alcohol at 70%..

In the same way, the place of injection must be also disinfected. Needles measuring 16 to 18 that have a usual length of 1 to 1 12 inches are the most suitable for the cattle. Injections administered through intramuscular route must be injected by introducing a needle having the proper size and length characteristics in the fleshy part of a thick muscle such as the haunches, hip of some parts of the muscle; avoid nerves. The treatment must be applied as soon as the first symptoms appears

AGRI-VETA LTD

Agriveta Spiramax

Spiramycin Injection 600 000 IU/ML

COMPOSITION: Each ml contains:

Spiramycin.....600 000 IU

Excipients .....1 ml

INDICATIONS:

Treatment of Acute clinical mastitis in lactating cows caused by Staphylococcus aureus strains sensitive to Spiramycin.
Treatment of Respiratory infections caused by Pasteurella multocida and Mannheimia haemolytica.

WITHDRAWAL PERIOD:

Mastitis: Meat and offal: 75 days. Milk: 13.5 days. Respiratory infections: Meat and offal: 75 days. Milk: In case of treatment at the dose required for respiratory diseases, the product is not authorised for use in animals producing milk for human consumption.

ADMINISTRATION AND DOSAGE:

Intramuscular route. Bodyweight should be determined as accurately as possible to avoid underdosing. Mastitis: 30,000 IU of Spiramycin per kg bodyweight (5 ml Spiramycin Inj. per 100 kg bodyweight) twice at 24h of interval. Respiratory infections: 100,000 IU of Spiramycin per kg bodyweight (5 ml Spiramycin Inj. per 30 kg bodyweight) twice at 48h of interval. Do not administer more than 15 ml per injection site. If the dose must be divided into two injections, then injections should be administered on opposite sides of the neck. If more than two injections are needed, a distance of at least 15 cm should be maintained between injections given on the same side of the neck. Failure to follow these instructions may result in residues above the established maximum residue

AGRI-VETA LTD

Agriveta OxyLong LA

brand of oxytetracycline

COMPOSITION: Each ml contains:

Oxytetracycline.....200 mg

Excipients .....1 ml

INDICATIONS:

Arthritis, gastrointestinal and respiratory infections caused by oxytetracycline sensitive microorganisms, like Bordetella, Campylobacter, Chlamydia, E. coli, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp. in calves, cattle, goa and sheep.

ADMINISTRATION AND DOSAGE:

For intramuscular administration: General: 1 ml per 10 kg body weight. This dosage can be repeated after 48 hours when necessary. Do not administer more than 20 ml in cattle and more than 5 ml in calves, goats and sheep per injection site.

ADMINISTRATION AND DOSAGE:

For intramuscular administration: General: 1 ml per 10 kg body weight. This dosage can be repeated after 48 hours when necessary. Do not administer more than 20 ml in cattle and more than 5 ml in calves, goats and sheep per injection site.

AGRI-VETA LTD

Agriveta TriCortin

0.1%+5%+10%+5%

COMPOSITION: Each ml contains:

Triamcinolone	1	mg
Hydrocortisone	50	mg
Phenyle Alanin	100	mg
Isoleucine	50	mg
Excipients	1 ml	

INDICATIONS:

This product comes with Triamcinolone acetonide and Hydrocortisone as active ingredient. Triamcinolone is an intermediate acting glucocorticoid and 4 - 10 times more potent than hydrocortisone. Intramuscular injection duration activity may persist up to 6 weeks. It is indicate in inflammatory or allergic disorders in dogs and cats like Feline inflammatory disease (IBD) disease (IBD) and also can be used in dermatologic disorders like dermatitis in cats.

ADMINISTRATION AND DOSAGE:

For intramuscular or subcutaneous injection. Dogs and cats 0.06 to 0.11 ml per kg body weight as a single injection in inflammatory or allergic disorders in dogs and cats and 0.11 ml per kg in dermatologic disorders residue

AGRI-VETA LTD

Agriveta Tyloxy Duo

Tylosin Tartrate + Oxytetracycline Injection 10% +10%

COMPOSITION: Each ml contains:

Tylosin tartrate.....100 mg
Oxytetracycline.....100 mg
Excipients .....1 ml

PHARMACOLOGICAL ACTION:

Tylosin acts bacteriostatically. It inhibits protein synthesis of susceptible microorganisms by binding to sub-units of the 50-S ribosome and by inhibition of the trans-location step. Tylosin has a wide spectrum of activity against gram-positive bacteria including Staphylococcus, Streptococcus, Coryne bacterium, and erysipelothrix. It has a much narrower gram-negative spectrum of activity, but has been shown to be active against campylobacter coli, and certain spirochaetes. It has also been shown to be extremely active against campylobacter coli, and certain spirochaetes. It has also been shown to be extremely active against Mycoplasma species isolated from both mammalian and avian hosts. Oxytetracycline is broad-spectrum antibacterial medicine, sensitive to rickettsia mycoplasma, chla-mydia, Spirochaeta. Other such as actinomycetes, bacillus anthracis, monocytosis listeria, clostridi-um, lave card bacteria genera, vibrio, gibraltar. Campylobacter, also has good effect on them.

PROPERTIES & INDICATIONS:

It has a broad antibacterial efficacy that includes Gram negative and positive bacteria, protozoas, siprochets, mycoplasmas and chlamydia. It is considered a first line treatment against metritis, mastitis, gastro-enteritis, infectious pneumonia and shipping fever, secondary bacterial infections and anaplasmosis and other infectious diseases.

AGRI-VETA LTD

Agriveta Tylosyn Plus

Tylosin Tratrate + Sulfadimidine + Trimethoprim Injection

COMPOSITION: Each ml contains:

Tylosin (as Tratrate).....100 mg
Sulfadimidine base.....125 mg
Trimethoprim.....25 mg
Excipients (ad.).....1 ml

DESCRIPTION:

The combination of Tylosin with Sulphadimidine and its enhancer Trimethoprim, complete its action and produces a distinctive effectiveness. Tylosin is an antibiotic substance with macrolide structure. It is high active against gram-positive bacteria and mycoplasma. Trimethoprim and sulphadimidine potentiate each other with bactericidal activity against many gram positive and gram negative bacteria

INDICATIONS:

For treatment of pneumoenteritis, foot infection and metritis in cattle and calves. Also for treatment of coli-bacillosis and pneumonia in sheep and goats, and bacterial scour in calves.

ADMINISTRATION AND DOSAGE:

By intramuscular injection: General dose: 1 ml/ 10 kg B.W. for 2-3 days. Cattle: 30-40 ml. Calves: 5-10 ml. Sheep: 5-10 ml. Lambs: 1-5 ml.

AGRI-VETA LTD

Agriveta TyloMax 200

Tylosin Tartrate Injection 20%

COMPOSITION: Each ml contains:
Tylosin tartrate.....200 mg
Excipients .....1 ml

INDICATIONS:
Gastrointestinal and respiratory infections causedby tylosin sensitive micro-organisms, likeCampylobacter, Mycoplasma, Pasteurella, Staphylococcus, Streptococcus and Treponema spp. in calves, cattle, goats and sheep.

ADMINISTRATION AND DOSAGE:
For intramuscular administration. General: 1 ml per 10-20 kg body weight for 3-5 days.

CONTRAINDICATIONS:
Hypersensitivity to tylosin. Concurrent administration of penicillins, cephalosporins, quinolones and cycloserine.

WITHDRAWAL PERIOD:
For meat: 10 days.
For milk: 3 days.

AGRI-VETA LTD

AZ TYLOMETHRIN PLUS

Tylosin Tratrate + Sulfadimidine + Trimethoprim Injection

COMPOSITION: Each ml contains:

Tylosin (as Tratrate).....100 mg
Sulfadimidine base.....125 mg
Trimethoprim.....25 mg
Excipients (ad.).....1 ml

INDICATIONS:

This medication is used to treat or prevent a lack of thiamine (deficiency) when the form taken by mouth cannot be used or would not work as well as the injection. Thiamine is a B vitamin that helps your body to use carbohydrates for energy.

ADMINISTRATION AND DOSAGE:

Dogs: 1 to 10 mg.
Cats: 0.5 to 5 mg.
Horses, Cattle: 100 mg Per 100 kg of body weight.
Administer intramuscularly, once daily or as recommended by the veterinarian.

WITHDRAWAL PERIOD:

Meat: Nil

AGRI-VETA LTD

Agriveta B-Complex Forte

Multivitamin Injection of Vit. B1, B2, B3, B5, B6, B12 & B15

COMPOSITION: Each ml contains:

Thiamine HCl (B1).....75.0 mg
Riboflavin (B2).....2.0 mg
Niacinamide (B3).....100.0 mg
D-Pantothenic (B5).....20.0 mg
Pyridoxine HCl (BG).....5.0 mg
Cyanocobalamin (B12).....100.0 mcg
Pangamic Acid (B15).....50.0 mg
Chlorobutanol.....0.5%
Benzyl Alcohol.....2.0%
Purified Water.....Q.S.

DESCRIPTION:

VITAMIN B COMPLEX is of value in convalescence from injury, surgery and illness and in geriatric conditions and parasitism.

INDICATIONS:

Maintains nerve functions, skin, hair, liver and muscle tone. Vital energy production system. Essential for red blood cell formation. Acts to reduce the effects of stress. Pangamic Acid (B15) Deoxident.

ADMINISTRATION AND DOSAGE:

Administer intramuscular or intravenous Dogs and cats: 0.5 to 2 ml: Sheep: 5 to 10 ml: Horses and cattle: 10 to 20 ml. Repeat daily as indicated.

AGRI-VETA LTD

Agriveta E-Sel Boost

Sodium Selenite + Vitamin E + VB. Injection

COMPOSITION: Each ml contains:

Sodium Selenite.....1 mg
Vitamin E.....60 mg
Vitamin B.....40 mg
Excipients (ad.).....1 ml

INDICATIONS:

Vitamin E deficiencies (like encephalomalacia, muscular dystrophy, exudative diathesis, decreased hatchability in eggs, infertility problems).

ADMINISTRATION AND DOSAGE:

For intramuscular administration: Calves, foals, goats and sheep: 2 ml/10 kg body weight, repeat after 2-3 weeks.

WITHDRAWAL PERIOD:

For meat: 28 days.

AGRI-VETA LTD

Agriveta GastroShield

Oral Antidiarrheal

For the treatment of gastrointestinal infections. Also for restoring electrolyte loss and eliminating bacterial toxins in camels, horses, cattle, goats, sheep, dogs & cats.

COMPOSITION: Each ml contains:

Sulphadimidine	21.3 mg
Sulphaguanidine	21.3 mg
Sulphadiazine	28.4 mg
Streptomycin sulphate	7.6 mg
Neomycin sulphate	1.8 mg
Hyoscine hydrobromide	0.02 mg
Sodium chloride	11.33 mg
Calcium gluconate	2.2 mg
Magnesium sulphate	0.6 mg
Potassium chloride	3.6 mg
Kaolin	103.3 mg
Pectin	7.1 mg
Glycine	20.9 mg

DOSE RATE:

For oral use only. First Dose 2 ml per 1 kg bodyweight. Following Doses 1 ml per 1 kg bodyweight night and morning for 3-5 days or for 24 hours after symptoms disappear.

AGRI-VETA LTD

Agriveta Calcivita

suspension

COMPOSITION: Each ml contains:

Calcium	56,000 mg
Phosphor	55,000 mg
Phosphate (Potassium)	10,000 mg
Potassium Gluconate	2,000 mg
Vitamin D3	100,000 IU
Iron	2,750 mg
Manganese	1,750 mg
Zinc	400 mg
Copper	70 mg
Cobalt	15 mg
Choline Chloride	2,000 mg
DL-methionine	1,000 mg
Lysine	10,000 mg
Citric Acid (mono)	1,000 mg
Purfified Water	Up to 1 L

INDICATION:

Improve performance of growth and fertility. In case of deficiencies in vitamins, minerals and trace element. When changing feeding habits. Help animal in the recovery during convalescence. In addition during antibiotic treatment. Larger resistance to infection. In addition during treatment or prevention of parasitic disease. Increase resistance under stress. Due to its high iron, vitamins and trace elements content, it helps the animal to combat anemia and to accelerate its recovery.

ADMINISTRATION AND DOSAGE:

Chicks/Broiler: 10ml/100 birds day for 5-7 days.
Growers: 25ml/100 birds day for 5-7 days.
Layers/Breeders: 50ml/100 birds day for 5-7 days. Cattle/Horse: 50ml/daily. Dogs: 0-10ml/dailys. Cat: 10 ml twice/daily.

AGRI-VETA LTD

Agriveta CyperFox 10

EXTERNAL PARASITICIDE

CONTAINS:

CYPERMETHRIN 10% W/V (100 g/L)

PROPERTIES & INDICATIONS:

CYPER-M-FOX: is Non-systemic insecticide with contact and stomach action for the control of and treatment of ectoparasites of livestock

DIRECTION FOR USE:

Cyper-M-Fox 10 formulation can affect fast on targeted pest. fox %10 EC: Veterinary Form/ EC Formulation with his unique SPECIES: Cattle, Camels, Goats, Sheep, etc.

INDICATIONS:

Helps control external parasitosis produced by flies, horse fly, ticks and louses in cattle and sheep inside sheds and our sheds also in close premises.

PESTS & RATE OF USE:

House on cattle and camells, 1ltr for 1000 ltr water -ticks on cattle and camels 1 ltr for 750 ltr water -ticks and louse on goats and sheep, 1 ltr for 800 ltr water -ticks and louse on chickens, 1 ltr for 750 ltr water (60 ltr solution for 1000 birds)

SITES:

Cattle, sheep, camels, hourses & poultry sheds out door & indoor premises

SAFETY PERIOD:

N/A

AGRI-VETA LTD

Agriveta DeltaSafe 2.5

Emulsifiable Concentrate

COMPOSITION: Each ml contains:

Deltamethrin (25g/L):.....2.5% (w/v)

Inert ingredient:.....97.5%

PROPERTIES OF PARKERS DELTA 2.5%:

A pyrethroid insecticide used on goats, sheep, camels and cows Can be used in several ways such as spraying or dipping. It is used to control external parasites such as fleas, lice and screwworms in goats, sheep, camels and cows. It is also used to control different types of war. No effectiveness in controlling flies and insects in stables, farms and barns

DESCRIPTION OF THE PEST:

A pyrethroid insecticide that is effective in eliminating many external parasites that infect various animals by spraying or dipping.

SAFETY PERIOD:

Czero for milk consumption and three days for meat consumption.

AGRI-VETA LTD

Agriveta HepaBest

EXTERNAL PARASITICIDE

COMPOSITION: Each ml contains:

Sorbitol	30,000 mg
Silymarin extract	3,000 mg
L-lysine	10,000 mg
Magnesium sulphate	50 mg
Choline chloride	30 mg
Methionine	5,000 mg
Nicotinamide	30,000 mg
Inositol	3,000 mg
Betain	8,000 mg
L-carnitine	4,000 mg
Aspartic acid	1,000 mg
Glutamic acid	5,000 mg
Artichoke	50,000 mg
Vitamin E	20 mg
Selenium	20 mg
Zinc sulphate	1200 mg

Liver is a vital organ involved in metabolism, storage, secretion, detoxification and protection. Being the central organ of metabolism, liver is always subject-ed to damages of various kinds. Liver stimulants and protectants are usually supplemented to support liver in detoxification and to improve the structural and functional integrity of liver cells. Hepa Best Oral solution contains all the essential nutrients formulated to protect, maintain and stimulate liver to perform various functions and to optimize productin.

Benefits Prevention of liver damage from mycotoxins & avoid perosis. Minimised liver damage due to pathogens like bacteria, virus, fungi & toovercome hypertrophy of liver. Increased liver secretions for better digetion and boost liver function. Faster recovery from stress & diseases of all types. Enhanced flock immunity,
Improved Growth, FCR, Ferdility & Hatchability.

AGRI-VETA LTD

Agriveta Levacloz

Levamisole HCL + Oxyclozanide Oral Suspension

COMPOSITION: Each ml contains:

Levamisole hydrochloride.....30 mg
Oxyclozanide.....60 mg
Excipients (ad.).....1 ml

PROPERTIES:

IOxyclozanide removes practically all flukes (Fasciola spp.) present in the bile ducts. Levamisole is the racemic mixture of optical isomers of tetramisole. Levamisole removes mature and developing immature stages of wide range of nematodes, and is highly effective against the following species: In the abomasum: Haemonchus, Ostertagia and Trichostrongylus spp. In the intestines: Trichostrongylus, Cooperia, Nematodirus, Oesophagostomum, Chabertia and Bunostomum. In the lungs: Dictyocaulus spp., including inhibited D. viviparus larvae.

INDICATIONS:

For the treatment and control of parasitic gastro-enteritis, verminous bronchitis, and liver flukes infestation. Gastro-intestinal worms: Nematodirus, Haemonchus, Ostertagia, Cooperia, Trichostrongylus, Bunostomum, Chabertia, Strongyloides, Oesophagostomum.
Lung worms: Dictyocaulus spp.
Tape worms: Moniezia spp. Liver flukes: Fasciola spp.

AGRI-VETA LTD

Agriveta PhoxiGuard

EXTERNAL PARASITICIDE

Each 1 L contain 50% Phoxim Properties & Indications:

PHOXIM 50% EC is an organophosphorous ectoparasiticide with high acaricidal efficacy and low toxicity to ward blooded animals. PHOXIM %50 EC is used for the treatment of infestations of pests, lice, fleas, External parasiticide crawling and flying insects (Dermanyssus gallinae) sensitive to organophosphates in the rearing buildings and in the presence of birds. PHOXIM %50 EC is highly toxic to fish and aquatic invertebrates.

WITHDRAWAL PERIOD:

For meat and offal consumption: 28 days from the last use. Eggs: 12 hours from the last treatment. Discard eggs laid on the same day of the treatment. For milk consumption: Not permitted in animals producing milk for human consumption. Antidote: Atropine sulphate. (Repeat after 30-15 minutes if necessar y)

STORAGE:

Store ectoparasiticide in its original containers tightly closed. - Keep away from children, food and drinks. - Store in dry, well ventilated place at temperature below 30 °C and away from direct sunlight.

RATE OF USE:

Dosage: Treatment of sheep, cattle and pigs is carried out by bathing, washing and spraying.

WASHING AND SPRAYING:

1.) Lice, fleas, flies, mites, and fly larvae once with a solution. of Phoxim per 10 liters of water.

2.) Scabies mites (Psoroptes, Sarcoptes, Chorioptes) twice with an interval of 7 days at the rate of 10 ml of Phoxim per 10 liters of water. In severe cases, the concentration of Phoxim is doubled 20 ml of Phoxim per 10 liters of water.

AGRI-VETA LTD

Agriveta DeltaPro 2.5

Used to control ticks on sheep, cow and goats

COMPOSITION EACH LT. CONTAINS:

Active Substance: Deltamethrin.....25 g

EMULSIFIERS:

Dodecylbenzene Sulphonate.....60 mg

Polyarylphenol ethoxylated.....40 mg

SOLVENT:

Solvesso.....up to 1 ml

FEATURES: DETERMENT 2.5% EC

A long-acting veterinary pesticide that works through contact and the digestive system.

- A powerful pesticide used to eliminate external parasites (ticks, lice, fleas, flies, and other insects). On sheep, goats, cows, poultry, camels and dogs.
- An easy-to-use pesticide that is stable under normal conditions of use.
-It does not evaporate and does not cause corrosion of the equipment.
- A veterinary pesticide used by spraying, washing or dipping.

SAFETY PERIOD:

Treated animal products may not be used until after 3 days.

MIXABILITY:

It can be mixed with most pesticides, except alkaline ones, and it is allowed to conduct a mini mixing trial 24 hours before use.

WARNING:

Avoid contaminating dairy products, milk, milking equipment, animal feed or water with the solution. Do not treat sick, tired, or exhausted animals.

AGRI-VETA LTD

Agriveta DeltaPro 5

Used to control ticks on sheep, cow and goats

COMPOSITION EACH 1ML CONTAINS:

50 mg Deltamethrin

PESTICIDE PROPERTIES:

Determent 5% EC insecticide in the form of an emulsifiable concentrate containing the active ingredient deltamethrin Mixed with water to combat ticks on cows, sheep and goats, by spraying or dipping the animals.

USES & RECOMMENDATION:

Animals Sheep, goats & cow Pests - ticks Dilution Rate 1Lt / 1920 Lt water Uses Dip and spray

NOTE:

Used only in animals that mention on table.

Safety Period Meat and Milk 2 days Always prepare a fresh solution when needed by adding concentrate to clean water only, stirring well. Do not store diluted product. Always wait 2 days minimum before batcher animals. Wait 2 days from the last treatment before used milk. Don't treat tired, sick, young (lower than 4 weeks), thirst & pregnancy animals. Don't used insecticide on injured animals.

AGRI-VETA LTD

Agriveta Herbalix

COMPOSITION ACTIVE INGREDIENT:

Cape Aloes.....9.70% w/w
Ammonium Acetate..... 60%..0.69% w/w
Peppermint Oil.....0.1% w/w

INDICATIONS:

For stomach upset. Not for Human Consumption.

ANIMALS SPECIES TO WHICH IT IS INDICATED:

Cattle, Camels, Sheep, Horses and Goats.

STORAGE:

Store In Shade And Dry Place. Temperature Below 30°C

AGRI-VETA LTD

Powder

Agriveta AmoxiClav Forte

TREATMENT OF RESPIRATORY AND DIGESTIVE TRACT INFECTIONS

INGREDIENTS:

Amoxicillin trihydrate.....16g
Clavulanate potassium.....4g
Excipient qs.....100g

INDICATIONS: Treatment of respiratory, digestive tract - infections, skin and soft tissue infections. - Use effective when cattle are mastitis, metritis, urinary tract infections, cystitis arthritis.

DOSAGE AND ADMINISTRATION: For via drinking water: Poultry: 1 g/2 liters of water. Treatment lasts for 3 to 5 days. Dogs, cats, calves, lambs: 0.5 gram per 1 Kg of live weight.

CONTRA INDICATIONS: Not to be administered to animals with known hypersensitivity to penicillins.

AGRI-VETA LTD

Powder

Agriveta BioPro Boost

Animal Feed Supplement

INGREDIENTS: Each kg contains:

Bacillus subtilis.....0.25 x 10 11CFU
Bacillus licheniformis.....0.25 x 10 11CFU
Bacillus pumilus.....0.25 x 10 11CFU
Lactobacillus acidophillus.....0.25 x 10 11CFU
Bacillus amyloliquefaciens.....0.2 x 10 10CFU
Total cellulase.....0.25 x 10 11CFU
Total Protease.....0.3 x 10 5U
Total Amylase.....0.3 x 10 50
VitaminA 2g
Vitamin D3.....0.3g

INDICATIONS: Improvement of nutrient absorption. Better balance of intestinal microflora. Improvement of Feed Conversion Ratio (FCR). - Prevention of digestion disease (diarrhea, constipation). - Improvement of feed efficiency and growth promotion, better, weight gain, egg production, fertility, hatchability.

ADMINISTRATION AND DOSAGE: Large Animals: Camel, horse, cattle: 10 gm/day. - Small Animals: Sheep, goat: 5 gm/day. Poultry: + Mix with feed: 1gm/2 kg of feed. + Dissolve in drinking water: 1gm/1-2 liters of drinking water, drink all day.

AGRI-VETA LTD

Agriveta CuSulfate

Copper Sulfate powder

INGREDIENTS: Each g contains:

Copper Sulfate.....800 mg

Excipients (ad.).....1g

TOPICAL USE: For wounds and ulcers: 0.1-5% antiseptic solutions. - Foot Bath: 5-10% solution twice daily for prevention and treatment of foot rot in sheep and cattle. Prevention of post-dipping lameness in sheep: 0.03% solution is added to the dipping bath.

INTERNAL USE: Cases of severe scouring in cattle due to excess molybdenum in pasture grass: Give daily 2 gm cow and 1 gm/ young animal for one week during grazing period as a drench or dietary supplement. Not given to calves while on milk or milk substitutes. Swayback prevention in lambs: The pregnants ewes in the affected areas are given copper sulphate, as a drench during gestation period: Per animal (1.5 gm) at 8 weeks. and then at (4) weeks before lambing. In copper deficient areas: add at rate of 0.5% to the salt. Feed additive as a production enhancer for dairy cattle and goats: Mix 250 gm /kg feed, then give a dose of (0.5-2 gm) from the mixture per animal

AGRI-VETA LTD

Agriveta QuadraSulfa

COMPOSITION: Each 1000 gm contains:

Sulfadiazine sodium.....9000 mg
Sulfathiazole sodium.....6000 mg
Sulfamirazine sodium.....15000 mg
Trimethoprim.....6000 mg

DESCRIPTION: This product has good antibacterial effects against both Gram positive and negative bacteria such as Streptococcus pyogenes, Salmonella, and Pneumonia. Sulfonamides are structurally similar to p- aminobenzoic acid and can compete with p-aminobenzoic acid for dihydrofolate synthase in bacteria, hindering the synthesis of dihydrofolate and ultimately affecting nucleic acid synthesis, inhibiting the action of sulfonamides can be antagonized by p-aminobenzoic acid and its derivatives (procaine, tetracaine). It has a good inhibitory effect on coccidia and Toxoplasma gondii.

INDICATIONS: Broad-spectrum antibacterial drugs. It is used to treat infectious diseases of livestock and poultry caused by sensitive bacteria such as gram-positive bacteria, gram-negative bacteria, mycoplasma, rickettsia, etc.

DOSAGE AND ADMINISTRATION: Poultry: 1 gm per 1 liter Withe Drinking Water for 3-5 Days Livestock: 0.25-0.5 gm per kg of body weight.

AGRI-VETA LTD

Agriveta Salic-C

INGREDIENTS:

Acetylsalicylic Acid (BP).....67mg
Vitamin C (BP).....200mg

INDICATIONS: Camel, Horse, Cattle, Sheep & Poultry: - PARKERS Sali-C Super powder is indicated for relief of pain, fever, pneumonia, inflammatory diseases, musculoskeletal disorders, heart diseases and stress due to vaccination, debeaking, transport, high humidity and temperature in poultry.

PHARMACOLOGICAL PROPERTIES: Acetylsalicylic Acid is readily absorbed from gastrointestinal tract and hydrolyzed into salicylic acid that inhibits the activity of cyclooxygenase and PGF2a. Salicylic acid also inhibits the synthesis of thromboxane, as a result platelet aggregation is inhibited and bleeding time prolongs. Salicylic acid has quick elimination and is mainly excreted as sodium salicylate in urine. Vitamin C participates in the oxido-reductive processes and aids in tissue respiration. It reduces the permeability of membranes, thickens the capillaries and helps to restrict the exudative inflammatory process. Vitamin C stabilizes adrenaline and assist the synthesis of corticosteroid hormones. Vitamin C also improves the appetite, secretion activity of gastric tract and the animal growth.

DOSAGE AND ADMINISTRATION:

Administer the following dose orally via drinking water. 1 gram of PARKERS Sali-C Super powder in 2 liters of drinking water.

AGRI-VETA LTD

Agriveta FosTylo

Best choice for gastrointestinal and respiratory tracts infections

COMPOSITION: Each 100 gr contains:

Fosfomycin.....20 gm
Tylosin Tarta.....5 gm
Dextro.....up to 100 gm

PROPERTIES & INDICATIONS: PARKERS TYLOPHOS contains a combination of broad spectrum antibiotics effective against a wide range of Gram positive and Gram negative bacteria and Mycoplasma. PARKERS TYLOPHOS is indicated for the treatment of gastrointestinal and respiratory tracts infections in poultry such as: pasteurellosis, salmonellosis, colibacillosis, CRD, infectious coryza, air sacculitis, infectious sinusitis, infectious synovitis, diarrhea, omphalitis and yolk sac infec-tions. PARKERS TYLOPHOS is indicated to relieve distress after vaccination, transporting and

DOSAGE & ADMINISTRATION: Poultry: 100 g/per 200 Liters of drinking water for 3-5 days.

WITHDRAWAL PERIOD: Treated poultry products should not be used for human consumption during treatment and within 7 days from the last dose.

AGRI-VETA LTD

Agriveta VitaGen 25

INGREDIENTS: Each 1 gram contains:

Paracetamol.....200 mg

Vitamin C.....50 mg

INDICATIONS: Parkers Vitagen 25% is used as antipyretic and - analgesic for animals small and large, in case of bacterial and viruses infections especially with gumboro in broilers and foot and mouth disease in sheep and cattle.

DOSAGE & ADMINISTRATION: Poultry: 200g/per 200 liters of water (For 3-5 days). Sheep and goats: 2-3 g Per day (For 3-5 days). Cows and camels: 4-6 g Per day (For 3-5 days). Dogs and Cats: 1-2 g Per day (For 3-5 days).

AGRI-VETA LTD

Agriveta Laxavit

Antiacid, Carminative, Laxative, Appetizer 101 gm.

INGREDIENTS: Each pack of 101g.contains:

Magnesium Carbonate:.....90 g
Sodium Carbonate:.....10 g
Nux Vomica:.....1g

Indications: Laxavit is indicated for the treatment of cases of - indigestion in animals e.g. acidosis, tympanitis constipation and abdominal poisoning caused by over feeding. It is recommended, as well as, appetizer, rumenal tonic and detoxicant.

Dosage: Cows, buffaloes, horses, camels: Dissolve contents of one pack
(101g) in one litre of drinking water and give as drench. Calves, sheep, goats: Dissolve (40g) of Laxavit in 0.5 litre of drinking water and give as drench.

Warning: Laxavit should not be given in cases of strychnine poisoning.

AGRI-VETA LTD



Agri Veta delivers veterinary medicines and vitamins built on quality, integrity, and customer-focused excellence.

AGRI-VETA LTD
RIEBECKSTRASSE 38 LEIPZIG PLZ 04317
DEUTSCHLAND,
GERMANY